

Acknowledgements

The authors express their sincere thanks to H. Y. Inoue and H. Matsueda, MRI, for their technical suggestions. The authors express their sincere thanks to K. Nemoto and for managing ^{85}Kr monitoring during the period from 1999 to 2001 and to M. Hirota, T. Tokieda, Y. Sawa and A. Wada for managing ^{85}Kr monitoring during the period from 2002 to 2005. M. Suzuki and S. Yasui of Shimadzu Corp., Y. Asai and I. Aoki of Repic Corp., and S. Yamamoto, KANSO Tecnos, are acknowledged for their services in constructing analytical system at MRI; without their expertise the present work could not have been accomplished. Y. Naka, Naka Showa, and T. Yogi and M. Yamada, Shimadzu Corp., are acknowledged for their services in purchasing various parts; without their expertise the present work also could not have been accomplished. Mr. S. Hodota, Nitto Koatsu, is acknowledged for devising a new plunger-type air pump. Mr. R. Lehnart, LND, is acknowledged for providing proportional counters which fulfilled our requirements. The authors express their sincere thanks to M. Suzuki, Shimadzu Corp., again for his great works to build new ^{85}Kr measurement system for JCAC in 2005 and 2006. K. Ito, Shimadzu Corp, is acknowledged for his technical support for JCAC. The authors also express their thanks to W.J.Zhu, S. Miyashita, and S. Ishikawa to prepare the whole text, figures and tables in this report.

This study is supported by the fund for the research of environmental radioactivity in Japan (Houshanou Chousa Kenkyuu-Hi) from the Ministry of Education, Culture, Sports, Science and Technology (MEXT), the former Science and Technology Agency (STA) throughout our study periods up to 2006.

References

- BfS, 2004: Krypton-Xenon sampling device, Manual for noble gas collection with the BfS system.
- Draxler, R. R., 2000: Demonstration of a global modeling methodology to determine the relative importance of local and long-distance sources. *Atmos. Environ.* **41**: 776-789.
- Fujitaka, K., 1995: External dose due to natural radiation in Japan. 'Human life with environmental radiation - collective effective dose from natural and man-made radiation sources', *Proceedings of the 22nd National Institute of radiological sciences seminar on environmental research, Chiba*, 1-2 Dec. 1994. Chiba: National Institute of Radiological Sciences (in Japanese).
- Geochemical Research Department, Meteorological Research Institute, 2005: Artificial Radionuclides in the Environment 2005. ISSN 1348-9739, Tsukuba, Japan
- Gurriaran, R., D. Maro, P. Bouisset, D. Hebert, G. Leclere, D. Mekhlouche, M. Rozet, L. Solier, 2004: In situ metrology of ^{85}Kr plumes released by the COGEMA La Hague nuclear reprocessing plant. *J. Environ. Radioact.*, **72**: 137-144.
- Hirota, M., K. Nemoto, A. Wada, Y. Igarashi, M. Aoyama, H. Matsuda, K. Hirose, H. Sartorius, C. Schlosser, S. Schmid, W. Weiss, K. Fujii, 2004: Spatial and Temporal Variations of Atmospheric ^{85}Kr Observed During 1995-2001 in Japan: Estimation of Atmospheric ^{85}Kr Inventory in the Northern Hemisphere. *J. Radiat. Res.*, **45**: 405-413.
- Igarashi, Y., M. Aoyama, T. Miyao, K. Hirose, K. Komura, and M. Yamamoto, 1999: Air concentration of radiocaesium in Tsukuba, Japan following the release from the Tokai waste treatment plant: comparisons of observations with predictions. *Appl. Radiat. Isot.*, **50**: 1063-1073.
- Igarashi, Y., M. Aoyama, K. Nemoto, K. Hirose, T. Miyao, K. Fushimi, M. Suzuki, S. Yasui, Y. Asai, I. Aoki, K. Fujii, S. Yamamoto, H. Sartorius, and W. Weiss, 2001: ^{85}Kr measurement system for continuous monitoring at Meteorological Research Institute, Japan. *J. Environ. Monit.*, **3**: 688-696.
- Igarashi, Y., T. Miyao, M. Aoyama, K. Hirose, H. Sartorius, and W. Weiss, 2000: Radioactive noble gases in surface air monitored at MRI, Tsukuba, before and after the JCO accident. *J. Environ. Radioact.*, **50**: 107-118.
- Igarashi, Y., H. Sartorius, T. Miyao, W. Weiss, K. Fushimi, M. Aoyama, K. Hirose, and H. Y. Inoue, 2000: ^{85}Kr and ^{133}Xe monitoring at MRI, Tsukuba and its importance. *J. Environ. Radioact.*, **48**: 191-202.
- Izrael, Y. A., I.M. Nazarov, A.G. Ryaboshapko, 1982: Release of man-made krypton-85 to the atmosphere. *Meteorologiya i Gidrologiya*, **6**: 5-15.

Jacob, D. J., M. J. Prather, S. C. Wofsy, and M. B. McElroy, 1987: Atmospheric distribution of ^{85}Kr simulated with a general circulation model. *Journal of Geophysical Research*, **92** (D6), 6614-6626.

JNC, 1999: JNC Technical Report, JNC TN8450 99 001. Japan Nuclear Cycle Development Institute (in Japanese).

JNC, 2002: JNC Technical Report, JNC TN8450 2002 002. Japan Nuclear Cycle Development Institute (in Japanese).

Kalinowski, M. B., H. Sartorius, S Uhl, W. Weiss, 2004: Conclusions on plutonium separation from atmospheric krypton-85 measured at various distances from the Karlsruhe reprocessing plant. *J. Environ. Radioact.*, **73**: 203-222

Loosli, H. H., 1992: Applications of ^{37}Ar , ^{39}Ar and ^{85}Kr in hydrology, oceanography, and atmospheric studies: Current state of the art, in *Isotopes of Noble Gases as Tracers in Environmental Studies; Proceedings of a Consultants Meeting*, International Atomic Energy Agency, Vienna. Pp.73-85.

MEXT, 2001: Annual Report on the Promotion of Science and Technology, 2001 (in Japanese)

NSC, 1997: Nuclear Safety Commission Monthly Report ISSN 0387-9674, 20 (5). Nuclear Safety Commission, Japan, 65-114 (in Japanese)

Pollard, D., V. Smith, D. Howett, E. Hayden, M. Fegan, M. O'Colmáin, J. D. Cunningham, 1997: Environmental radioactivity surveillance programme 1994-1996. Radiological Protection Institute of Ireland.

Rath, H. K., 1988: Simulation der globalen ^{85}Kr und $^{14}\text{CO}_2$ Verteilung mit Hilfe eines zeitabhängigen, zweidimensionalen Modells der Atmosphäre. Dissertation, Universität Heidelberg.

Sartorius, H.; C. Schlosser, S. Schmid, W. Weiss, (2002: Verfahren zur Bestimmung der Aktivitätskonzentration der atmosphärischen Edelgase Krypton-85 und Xenon-133, Empfehlungen zur Überwachung der Umweltradioaktivität, Loseblattsammlung FS-788-15-AKU, Blatt 3.4.9, Juli 2002-11-06, http://osiris2.pi-consult.de/userdata/1_2/p_25/library/data/lb3-4-9_2.pdf. (in German)

Sittkus A. and H. Stockburger, 1976: Krypton-85 als Indikator des Kernbrennstoffverbrauchs. Naturwissenschaften 63, 266-272. (in German)

STA, 1997: On the investigation into the cause of a fire and explosion accident at the bituminisation facility, Power Reactor and Fuel Development, Tokai Works. Nuclear Safety Bureau, Science and Technology Agency, Japan (in Japanese).

Stockburger, H., H. Sartorius, and A. Sittkus, 1977: Messung der Krypton-85 und Xenon-133 Aktivität der atmosphärischen Luft. *Zeitschrift für Naturforschung*, **32**:

1249-1253. (in German)

Styra, B. and D. Butkus, 1991: Geophysical Problems of Krypton.85 in the Atmosphere, Hemisphere, New York/Mokslas, Vilnius.

United Nations, 2000: Sources and Effects of Ionizing Radiation, United Nations Scientific Committee on the Effects of Atomic Radiation, UNSCEAR 2000 Report to the General Assembly, with Scientific Annexes. United Nations Sales Publication E.00.IX.3, Vol. 1, United Nations, New York, 280-281.

Von Hippel, F., D. H. Albright, G. P. Levi, 1986: Quantities of Fissile Materials in the US and Soviet Nuclear Weapon Arsenals, Rep. PU/CEES 168, Center for Energy and Environmental Studies, Protection University, New Jersey.

Weiss, W. H. Stockburger, H. Sartorius, K. Rozanski, C. Heras, H.G. Östlund, 1986: Mesoscale Transport of ^{85}Kr Originating from European Sources. NIM B17, 571-574.

Weiss, W., H. Sartorius, and H. Stockburger, 1992: Global distribution of atmospheric ^{85}Kr . A database for the verification of transport and mixing models. In Isotopes of noble gases as tracers in environmental studies. IAEA, Vienna, 29-62

Weiss, W., A. Sittkus, H. Sartorius, and Stockburger, H., 1987: Tschernobyl . eine Rückschau. *Phys. Bl.*, **43** (5), 125-130.

World Meteorological Organization (WMO), 1996: Report of an Expert Consultation on ^{85}Kr and ^{222}Rn : Measurements, Effects and Applications (Freiburg, Germany, 28-31 March 1995), Environmental Pollution Monitoring and Research Programme Report Series No. 109.

Zimmermann, P. H., J. Feichter, H. K. Rath, P. J. Crutzen, and W. Weiss, 1989: A global three-dimensional source-receptor model investigation using ^{85}Kr . *Atmospheric Environment*, **23**, 25-35.