

Acknowledgement

We appreciate the generous technical support from members of the numerical forecast group of the Climate Prediction Division of the JMA. Mr. Noriyuki Adachi of the JMA helped us greatly with the implementation of the operational BGM cycle system of the JMA onto the MRI supercomputing system. Mr. Hitoshi Sato of JMA gave us much useful advice about the operational BGM cycle system of the JMA.

References

- Adachi, Y., S. Yukimoto, M. Deushi, A. Obata, H. Nakano, T. Y. Tanaka, M. Hosaka, T. Sakami, H. Yoshimura, M. Hirabara, E. Shindo, H. Tsujino, R. Mizuta, S. Yabu, T. Koshiro, T. Ose, and A. Kitoh, 2013: Basic performance of a new earth system model of the Meteorological Research Institute (MRI-ESM1). *Pap. Meteor. Geophys.*, **64**, 1–19.
- Buizza, R., M. J. Miller, and T. N. Palmer, 1999: Stochastic simulation of model uncertainties in the ECMWF ensemble prediction system. *Q. J. R. Meteorol. Soc.*, **125**, 2887–2908.
- Buizza, R., Tribbia, J., Molteni, F., and Palmer, T. N., 1993: Computation of optimal unstable structures for a numerical weather prediction model. *Tellus*, **45A**, 388–407.
- Chikamoto, Y., H. Mukougawa, T. Kubota, H. Sato, A. Ito, and S. Maeda, 2007: Evidence of growing bred vector associated with the tropical intraseasonal oscillation, *Geophys. Res. Lett.*, **34**, L04806, doi:10.1029/2006GL028450.
- Dee D.P. et al. 2011: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. R. Meteorol. Soc.*, **137**, 553–597, doi:10.1002/qj.828.
- Endo, H., A. Kitoh, T. Ose, R. Mizuta, and S. Kusunoki, 2012: Future changes in Asian precipitation simulated by multi-physics and multi-sea surface temperature ensemble experiments with high-resolution Meteorological Research Institute atmospheric general circulation models (MRI-AGCMs). *J. Geophys. Res.*, **117**, D16118, doi:10.1029/2012JD017874.
- Hirai, M., T. Sakashita, H. Kitagawa, T. Tsuyuki, M. Hosaka, and M. Oh'izumi, 2007: Development

and validation of a new land surface model for JMA's operational global model using the CEOP observation dataset. *J. Meteor. Soc. Japan*, **85A**, 1–24.

Ishii, M., A. Shouji, S. Sugimoto and T. Matsumoto, 2005: Objective Analyses of Sea-Surface Temperature and Marine Meteorological Variables for the 20th Century Using ICOADS and the KOBE Collection. *Int. J. of Climatology*, **25**, 865-879.

Kawai, H., 2006: PDF cloud scheme and prognostic cloud scheme in JMA global model. *CAS/JSC WGNE Research Activities in Atmospheric and Ocean Modeling*, **36**, 4.15–4.16.

Kyoda, M., 2000: Performance of weekly ensemble forecast system. NPD/JMA Technical Report, **47**, Numerical Prediction Division, Japan Meteorological Agency. ,86-93. (in Japanese)

Kyoda, M., 2006: Weekly ensemble forecast. NPD/JMA Technical Report, **52**, Numerical Prediction Division, Japan Meteorological Agency. ,23-33. (in Japanese)

Legras, B. and R. Vautard, 1995: A guide to Liapunov vectors. Proceedings of the ECMWF seminar on Predictability, 143-156.

Lorenz, E. N., 1963: Deterministic non-periodic flows. *J. Atmos. Sci.*, **20**, 130–141.

Mizuta, R., H. Yoshimura, H. Murakami, M. Matsueda, H. Endo, T. Ose, K. Kamiguchi, M. Hosaka, M. Sugi, S. Yukimoto, S. Kusunoki, and A. Kitoh, 2012: Climate simulations using MRI-AGCM3.2 with 20-km grid. *J. Meteor. Soc. Japan*, **90A**, 233-258.

Mukougawa H., H. Sakai, and T. Hirooka, 2005: High sensitivity to the initial condition for the prediction of stratospheric sudden warming. *Geophys. Res. Lett.*, **32**, L17806, doi:10.1029/2005GL022920.

Murakami, H., and T. Matsumura, 2007: Development of an effective non-linear normal mode initialization method for a high resolution global model. *J. Meteor. Soc. Japan*, **85**, 187-208.

Murakami, H., Y. Wang, H. Yoshimura, R. Mizuta, M. Sugi, E. Shindo, Y. Adachi, S. Yukimoto, M. Hosaka, S. Kusunoki, T. Ose, and A. Kitoh, 2011: Future changes in tropical cyclone activity projected by the new high-resolution MRI-AGCM. *J. Climate*, **25**, 3237-3260.

Onogi, K., J.Tsutsui, H.Koide, M. Sakamoto, S. Kobayashi, H. Hatsushika, T. Matsumoto, N. Yamazaki, H. Kamahori, K. Takahashi, S. Kadokura, K. Wada, K. Kato, R. Oyama, T. Ose, N. Mannoji and R. Taira, 2007: The JRA-25 Reanalysis. *J. Meteor. Soc. Japan*, **85**, 369–432.

Sellers, P. J., Y. Mints, Y. C. Sud and A. Dalcher, 1986: A simple biosphere model (SiB) for use within general circulation models. *J. Atmos. Sci.*, **43**, 505–531.

Taylor, K. E., R. J. Stouffer, and G. A. Meehl, 2012: An overview of CMIP5 and the experiment design, *Bull. Am. Meteorol. Soc.*, **93**, 485–498, doi:10.1175/BAMS-D-11-00094.1.

Tiedtke, M., 1993: Representation of clouds in largescale models. *Mon. Wea. Rev.*, **121**, 3040–3061.

Toth, Z., and E. Kalnay, 1993: Ensemble forecasting at NMC: The generation of perturbations. *Bull. Amer. Meteor. Soc.*, **74**, 2317–2330.

Toth, Z., and E. Kalnay, 1997: Ensemble forecasting at NCEP and the breeding method. *Mon. Wea. Rev.*, **125**, 3297–3319.

Yamaguchi, M., R. Sakai, M. Kyoda, T. Komori, and T. Kadokawa, 2009: Typhoon Ensemble Prediction System developed at the Japan Meteorological Agency. *Mon. Wea. Rev.*, **137**, 2592–2604.

Yukimoto, S., H. Yoshimura, M. Hosaka, T. Sakami, H. Tsujino, M. Hirabara, T. Y. Tanaka, M. Deushi, A. Obata, H. Nakano, Y. Adachi, E. Shindo, S. Yabu, T. Ose and A. Kitoh, 2011: Meteorological Research Institute-Earth System Model v1 (MRIESM1) --Model description--. *Tech. Rep. Meteor. Res. Inst.*, **64**, 88 pp.