

References

- Aoki, S., T. Nakazawa, S. Murayama, and S. Kawaguchi (1992), Measurements of atmospheric methane at the Japanese Antarctic, *Syowa, Tellus, Ser. B*, **44**, 273-281.
- Slugokenky, E. J., R. C. Myers, P. M. Lang, K. A. Masarie, A. M. Crotwell, K. W. Thoning, B. D. Hall, J. W. Elkins, and L. P. Steele (2005), Conversion of NOAA atmospheric dry air CH₄ mole fractions to a gravimetrically prepared standard scale, *J. Geophys. Res.*, **110**, D18306, doi:10.1029/2005JD006035.
- Flores, E., J. Viallon, T. Choteau, P. Moussay, R. I. Wielgosz, N. Kang, B. M. Kim, E. Zalewska, A. van der Veen, L. Konopelko, H. Wu, Q. Han, G. Rhoderick, F. R. Guenther, T. Watanabe, T. Shimosaka, K. Kato, B. Hall, and P. Brewer (2015), International comparison CCQM-K82: methane in air at ambient level (1800 to 2200) nmol/mol, *Metrologia*, **52**, 08001, doi:10.1088/0026-1394/52/1A/08001.
- Hall, B. D., G. S. Dutto, and J. W. Elkins (2007), The NOAA nitrous oxide standard scale for atmospheric observations, *J. Geophys. Res.*, **112**, D09305, doi:10.1029/2006JD007954.
- ISO 6142:2001, “Gas analysis - Preparation of calibration gas mixtures - Gravimetric method”, second edition.
- Katsumata, K., T. Machida, H. Tanimoto, H. Nara, and H. Mukai (2011), Re-evaluation of NIES CO scale using high concentration gravimetric CO standard gases, *GAW Report*, **194**, 295-298.
- Kawasaki, T., A. Takizawa, M. Takahashi, H. Koide, T. Nakazawa, S. Aoki, S. Morimoto, K. Kato, T. Shimosaka, N. Aoki, T. Watanabe, T. Machida, Y. Tohjima, H. Mukai, K.

- Katsumata, T. Fujitani, S. Murayama, S. Ishidoya, D. Goto, H. Matsueda, Y. Sawa, and K. Tsuboi (2016), Inter-comparison experiments of standard gases for JMA/WCC activity, *18th WMO/IAEA Meeting on Carbon Dioxide, Other Greenhouse Gases and Related Tracers Measurement Techniques (GGMT-2015)*, GAW Rep. 229, pp. 104-109, World Meteorol. Organ., Geneva, Switzerland.
- Machida, T., T. Nakazawa, Y. Fujii, S. Aoki, and O. Watanabe (1995), Increase in the atmospheric nitrous oxide concentration during the last 250 years, *Geophys. Res. Lett.*, **22**, 2921-2924.
- Machida, T., H. Matsueda, Y. Sawa, Y. Nakagawa, K. Hirotani, N. Kondo, K. Goto, T. Nakazawa, K. Ishikawa, and T. Ogawa (2008), Worldwide measurements of atmospheric CO₂ and other trace gas species using commercial airlines, *J. Atmos. Oceanic Technol.*, **25**, 1744-1754, doi:10.1175/2008JTECHA1082.1.
- Machida, T., Y. Tohjima, K. Katsumata, and H. Mukai (2011), A new CO₂ calibration scale based on gravimetric one-step dilution cylinders, in *15th WMO/IAEA Meeting of Experts on Carbon Dioxide, Other Greenhouse Gases and Related Tracers Measurement Techniques*, National Institute for Environmental Studies, GAW Rep. 194, pp. 114-119, World Meteorol. Organ., Geneva, Switzerland.
- Matsueda, H. (1993), Intercalibration experiment of methane standard gas scale between NOAA/CMDL and MRI/GRL, *Pap. Meteorol. Geophys.*, **44**, 45-56.
- Matsueda, H., and H. Y. Inoue (1996), Measurements of atmospheric CO₂ and CH₄ using a commercial airliner from 1993 to 1994, *Atmos. Environ.*, **30**, 1647-1655.

Matsueda, H., H.Y. Inoue, Y. Sawa, Y. Tsutsumi, and M. Ishii (1998), Carbon monoxide in the upper troposphere over the western Pacific between 1993 and 1996, *J. Geophys. Res.*, **103**, 19093-19110.

Matsueda, H., Y. Sawa, A. Wada, H. Y. Inoue, K. Suda, Y. Hirano, K. Tsuboi, and S. Nishioka (2004a), Methane standard gases for atmospheric measurements at the MRI and JMA and intercomparison experiments, *Pap. Meteorol. Geophys.*, **54**, 91-109, doi:10.2467/mripapers.54.91.

Matsueda, H., K. Suda, S. Nishioka, Y. Hirano, Y. Sawa, K. Tsuboi, Y. Tutumi, H. Kamiya, K. Nemoto, H. Nagai, M. Yoshida, S. Iwano, O. Yamamoto, H. Morishita, M. Kamata, and A. Wada (2004b), Re-evaluation for scale and stability of CO₂ standard gases used as long-term observations at the Japan Meteorological Agency and the Meteorological Research Institute (in Japanese), *Technical Reports of Meteorological Research Institute*, **45**, 1-38.

Murayama, S., N. Saigusa, D. Chan, S. Yamamoto, H. Kondo and Y. Eguchi (2003), Temporal variations of atmospheric CO₂ concentration in a temperate deciduous forest in central Japan, *Tellus, Ser. B*, **55**, 232-243.

Murayama, S., C. Takamura, S. Yamamoto, N. Saigusa, S. Morimoto, H. Kondo, T. Nakazawa, S. Aoki, T. Usami, and M. Kondo (2010), Seasonal variations of atmospheric CO₂, δ¹³C, and δ¹⁸O at a cool temperate deciduous forest in Japan: Influence of Asian monsoon, *J. Geophys. Res.*, **115**, D17304, doi:10.1029/2009JD013626.

Novelli, P. C., K. A. Masarie, P. M. Lang, B. D. Hall, R. C. Myers, and J. W. Elkins (2003),

Reanalysis of tropospheric CO trends: Effects of the 1997–1998 wildfires, *J. Geophys. Res.*, **108**, D15, 4464, doi:10.1029/2002JD003031.

Novelli, P. C. (2016), Round robin of carbon monoxide measurements among Japan, Switzerland and the USA (unpublished report available upon request).

Takahashi, M., T. Nakazawa, S. Aoki, D. Goto, K. Kato, N. Aoki, T. Watanabe, T. Machida, Y. Tohjima, K. Katsumata, S. Murayama, S. Ishidoya, S. Morimoto, T. Fujitani, H. Koide, A. Takizawa, H. Matsueda, Y. Sawa, and K. Tsuboi (2013), Intercomparison experiments for greenhouse gases observation (iceGGO) in Japan., *Asian-Pacific GAW Greenhouse Gases Newsletter*, **4**, 45-49.

Takahashi, M., T. Nakazawa, S. Aoki, D. Goto, K. Kato, N. Aoki, T. Watanabe, T. Machida, Y. Tohjima, K. Katsumata, S. Murayama, S. Ishidoya, S. Morimoto, T. Fujitani, H. Koide, A. Takizawa, H. Matsueda, Y. Sawa, and K. Tsuboi (2014), Intercomparison experiments for greenhouse gases observation (iceGGO) in Japan, *17th WMO/IAEA Meeting on Carbon Dioxide, Other Greenhouse Gases and Related Tracers Measurement Techniques (GGMT-2013)*, WMO/GAW Report, No. **213**, 138-143.

Tanaka, S., T. Nakazawa, and S. Aoki (1983), High quality measurements of the concentration of atmospheric carbon dioxide, *J. Meteorol. Soc. Japan*, **61**, 678-685.

Tohjima, Y., K. Katsumata, I. Morino, H. Mukai, T. Machida, I. Akama, T. Amari, and U. Tsunogai (2009), Theoretical and experimental evaluation of the isotope effect of NDIR analyzer on atmospheric CO₂ measurement, *J. Geophys. Res.*, **114**, D13302, doi:10.1029/2009JD011734.

- Tsuboi, K., H. Matsueda, Y. Sawa, Y. Niwa, M. Nakamura, D. Kuboike, K. Saito, H. Ohmori, S. Iwatsubo, H. Nishi, Y. Hanamiya, K. Tsuji, and Y. Baba (2013), Evaluation of a new JMA aircraft flask sampling system and laboratory trace gas analysis system, *Atmos. Meas. Tech.*, **6**, 1257-1270, doi:10.5194/amt-6-1257-2013.
- Tsuboi, K., H. Matsueda, Y. Sawa, Y. Niwa, M. Takahashi, S. Takatsuji, T. Kawasaki, T. Shimosaka, T. Watanabe, and K. Kato (2016), Scale and stability of methane standard gas in JMA and comparison with MRI standard gas, *Pap. Meteorol. Geophys.*, **66**, 15-24, doi:10.2467/mripapers66.15.
- WMO (2016a), WMO Greenhouse Gas Bulletin: The state of greenhouse gases in the atmosphere based on global observations through 2015, No.12.
- WMO (2016b), 18th WMO/IAEA Meeting on Carbon Dioxide, Other Greenhouse Gases and Related Tracers Measurement Techniques (GGMT-2015), GAW Report, No. 229.
- Yashiro, H., S. Sugawara, K. Sudo, S. Aoki, and T. Nakazawa (2009), Temporal and spatial variations of carbon monoxide in the western part of the Pacific Ocean, *J. Geophys. Res.*, **114**, D08305, doi:10.1029/2008JD010876.
- Zhao, C. L., and P. P. Tans (2006), Estimating uncertainty of the WMO mole fraction scale for carbon dioxide in air, *J. Geophys. Res.*, **111**, D08S09, doi:10.1029/2005JD006003.