

参考文献

- Bloom, S. C. , L. L. Takacs, Silva A. M. D., and Ledvina D., 1996: Data assimilation using incremental analysis updates. *Mon. Wea. Rev.*, **124**, 1256-1271.
- Boyer, T. P., J. I. Antonov, O. K. Baranova, C. Coleman, H. E. Garcia, A. Grodsky, D. R. Johnson, R. A. Locarnini, and M. M. Zweng, 2013: World Ocean Database 2013. *NOAA Atlas NESDIS* **72**.
- Byun, D. S., and D. E. Hart, 2017: A robust interpolation procedure for producing tidal current ellipse inputs for regional and coastal ocean numerical models. *Ocean Dyn.*, **67**, 451-463. <https://doi.org/10.1007/s10236-017-1037-4>
- Fujii, Y., and M. Kamachi, 2003: Three-dimensional analysis of temperature and salinity in the Equatorial Pacific using a variational method with vertical coupled temperature-salinity EOF modes. *J. Geophys. Res.*, **108**, 3297. <https://doi.org/10.1029/2002JC001745>
- Hamilton, D., 1994: GTSPP builds an ocean temperature-salinity database. *Earth Syst. Monitor*, **4**, 4-5.
- Hirose, N., N. Usui, K. Sakamoto, H. Tsujino, G. Yamanaka, H. Nakano, S. Urakawa, T. Toyoda, Y. Fujii, and N. Kohno, 2019: Development of a new operational system for monitoring and forecasting coastal and open ocean states around Japan. *Ocean Dyn.*, **69**, 1333-1357. <https://doi:10.1007/s10236-019-01306-x>
- Hunke, E. C., and J. K. Dukowicz, 1997: An elastic-viscous-plastic model for sea ice dynamics. *J. Phys. Oceanogr.*, **27**, 1849-1867. [https://doi.org/10.1175/1520-0485\(1997\)027<1849:AEVPMF>2.0.CO;2](https://doi.org/10.1175/1520-0485(1997)027<1849:AEVPMF>2.0.CO;2)
- Hunke, E. C., and J. K. Dukowicz, 2002: The elastic-viscous-plastic sea ice dynamics model in general orthogonal curvilinear coordinates on a sphere: incorporation of metric terms. *Mon. Wea. Rev.*, **130**, 1848-1865. [https://doi.org/10.1175/1520-0493\(2002\)130<1848:TEVPSI>2.0.CO;2](https://doi.org/10.1175/1520-0493(2002)130<1848:TEVPSI>2.0.CO;2)
- Kobayashi, S., Y. Ota, Y. Harada, A. Ebita, M. Moriya, H. Onoda, K. Onogi, H. Kamahori, C. Kobayashi, H. Endo, K. Miyaoka, and K. Takahashi, 2015: The JRA-55 reanalysis: general specifications and basic characteristics. *J. Meteor. Soc. Japan*, **93**, 5–48. <https://doi.org/10.2151/jmsj.2015-001>
- 栗原幸雄, 桜井敏之, 倉賀野連, 2006: 衛星マイクロ波放射計, 衛星赤外放射計及び現場観測データを用いた全球日別海面水温解析 . 測候時報, **73**, S1-S18 .
- Large, W. G., and S. G. Yeager, 2004: Diurnal to decadal global forcing for ocean and sea-ice models: The data sets and flux climatologies. *NCAR Tech. Note*, TN-460+STR, CGD Division of the National Center for Atmospheric Research.
- Mellor, G. L., and L. Kantha, 1989: An ice-ocean coupled model. *J. Geophys. Res.*, **94**, 10937-10954. <https://doi.org/10.1029/JC094iC08p10937>
- Sakamoto, K., H. Tsujino, H. Nakano, M. Hirabara, and G. Yamanaka, 2013: A practical scheme to introduce explicit tidal forcing into an OGCM. *Ocean Sci.*, **9**, 1089-1108. <https://doi.org/10.5194/os-9-1089-2013>
- Sakamoto, K., H. Tsujino, H. Nakano, S. Urakawa, T. Toyoda, N. Hirose, N. Usui, and G. Yamanaka, 2019: Development of a 2km-resolution ocean model covering the coastal seas around Japan for operational application. *Ocean Dyn.*, **69**, 1181-1202. <https://doi.org/10.1007/s10236-019-01291-1>
- Suzuki, T., D. Yamazaki, H. Tsujino, Y. Komuro, H. Nakano, and S. Urakawa, 2018: A dataset of con-

- tinental river discharge based on JRA-55 for use in a global ocean circulation model. *J. Oceanogr.*, **74**, 421-429. <https://doi.org/10.1007/s10872-017-0458-5>
- Tsujino, H., H. Nakano, K. Sakamoto, S. Urakawa, M. Hirabara, H. Ishizaki, and G. Yamanaka, 2017: Reference manual for the Meteorological Research Institute Community Ocean Model version 4 (MRI.COMv4). *Tech. Rep.*, **80**, Meteorological Research Institute, Japan. <https://doi.org/10.11483/mritechrepo./80>
- Tsujino, H., S. Urakawa, H. Nakano, R. J. Small, W. M. Kim, S. G. Yeager, G. Danabasoglu, T. Suzuki, J. L. Bamber, M. Bentsen, C. W. Boning, A. Bozec, E. P. Chassignet, E. Curchitser, F. B. Dias, P. J. Durack, S. M. Griffies, Y. Harada, M. Ilicak, S. A. Josey, C. Kobayashi, S. Kobayashi, Y. Komuro, W. G. Large, J. L. Sommer, S. J. Marsland, S. Masina, M. Scheinert, H. Tomita, M. Valdivieso, and D. Yamazaki, 2018: JRA-55 based surface dataset for driving ocean-sea-ice models (JRA55-do). *Ocean Modell.*, **130**, 79-139. <https://doi.org/10.1016/j.ocemod.2018.07.002>
- 碓氷典久, 今泉孝男, 辻野博之, 2010: MOVE/MRI.COM への海水密度同化の導入に向けて - オホーツク海を対象とした予備調査と簡易同化実験 - . *測候時報*, **77**, s72-s82.
- Usui, N., Y. Fujii, K. Sakamoto, M. Kamachi, 2015: Development of a four dimensional variational assimilation system toward coastal data assimilation around Japan. *Mon. Wea. Rev.*, **143** 3874-3892. <https://doi.org/10.1175/MWR-D-14-00326.1>
- Usui, N., T. Wakamatsu, Y. Tanaka, N. Hirose, T. Toyoda, S. Nishikawa, Y. Fujii, Y. Takatsuki, H. Igarashi, H. Nishikawa, Y. Ishikawa, T. Kuragano, and M. Kamachi, 2017: Four-dimensional variational ocean reanalysis: a 30-year high-resolution dataset in the western North Pacific (FORA-WNP30). *J. Oceanogr.*, **73**, 205-233. <https://doi.org/10.1007/s10872-016-0398-5>