

H. References

- Adachi, K., M. Kajino, Y. Zaizen, and Y. Igarashi, 2013: Emission of spherical cesium-bearing particles from an early stage of the Fukushima nuclear accident. *Nature Scientific Reports*, **3**, 2554, doi:10.1038/srep02554.
- Araki, K., 2000: Six-hour forecasts of precipitation. *Reports of the Numerical Prediction Division*, **47**, 36–41. (in Japanese)
- Arnold, D., C. Maurer, G. Wotawa, R. Draxler, K. Saito, and P. Seibert, 2015: Influence of the meteorological input on the atmospheric transport modelling with FLEXPART of radionuclides from the Fukushima Daiichi nuclear accident. *J. Environ. Radioact.*, **139**, 212–225.
- Becker, A., G. Wotawa, L-E De Geer, P. Seibert, R. Draxler, C. Sloan, R. D'Amours, M. Hort, H. Glaab, P. Heinrich, Y. Grillon, V. Shershakov, K. Katayama, Y. Zhang, P. Stewart, M. Hirtl, M. Jane, and P. Chen, 2007: Global backtracking of anthropogenic radionuclides by means of a receptor oriented ensemble dispersion modelling system in support of Nuclear-Test-Ban Treaty verification. *Atmos. Environ.*, **41**, 4520–4534.
- Bedwell, P., J. Wellings, S. M. Haywood, A. R. Jones, and M. C. Hort, 2011: Cloud gamma modelling in the UK Met Office's NAME III model. *AGU Chapman Conference*.
- Blackadar, A. K., 1962: The vertical distribution of wind and turbulent exchange in a neutral atmosphere. *J. Geophys. Res.*, **67**, 3095–3102.
- Browning, K. A., and F. F. Hill, 1981: Orographic Rain. *Weather*, **35**, 326–329.
- Burgin, L. E., J. Gloster, C. Sanders, P. S. Mellor, S. Gubbins, and S. Carpenter, 2012: Investigating incursions of bluetongue virus using a model of long-distance culicoides biting midge dispersal. *Transboundary and Emerging Diseases*, doi:10.1111/j.1865-1682.2012.01345.x.
- Chino, M., H. Nakayama, H. Nagai, H. Terada, G. Katata, and H. Yamazawa, 2011: Preliminary estimation of release amounts of ^{131}I and ^{137}Cs accidentally discharged from the Fukushima Daiichi Nuclear Power Plant into the atmosphere. *J. Nucl. Sci. Technol.*, **48**, 1129–1134.
- Christoudias, T., and J. Lelieveld, 2013: Modelling the global atmospheric transport and deposition of radionuclides from the Fukushima Dai-ichi nuclear accident. *Atmos. Chem. Phys.*, **13**, 1425–1438.
- Clark, M. J., and F. B. Smith, 1988: Wet and dry deposition of Chernobyl releases. *Nature*, **332**, 245–249.
- Courtier, P., J.-N. Thépaut, and A. Hollingsworth, 1994: A strategy for operational implementation of 4D-Var, using an incremental approach. *Q. J. Roy. Met. Soc.*, **120**, 1367–1387.
- D'Amours, R., and A. Malo, 2004: A zeroth order Lagrangian particle dispersion model MLDPO. *Internal publication*, Canadian Meteorological Centre, Environmental Emergency Response Section, Dorval, QC, Canada, 19 pp. (available online at <http://eer.cmc.ec.gc.ca/publications/DAmours.Malo.2004.CMC-EER.MLDPO.pdf>)
- D'Amours, R., A. Malo, R. Servranckx, D. Bensimon, S. Trudel, and J.-P. Gauthier, 2010: Application of the atmospheric Lagrangian particle dispersion model MLDPO to the 2008 eruptions of Okmok

- and Kasatochi volcanoes. *J. Geophys. Res.*, **115**, D00L11, 11 pp, doi:10.1029/2009JD013602.
- Delage, Y., 1997: Parameterising sub-grid scale vertical transport in atmospheric models under statically stable conditions. *Bound.-Layer Meteor.*, **82**, 23–48.
- Draxler, R. R., 1999: HYSPLIT_4 User's Guide. *NOAA Tech. Memo, ERL ARL-230, Air Resources Laboratory, Silver Spring, MD*, 35 pp. (available online at http://www.arl.noaa.gov/documents/reports/hysplit_user_guide.pdf)
- Draxler, R. R., 2006: The Use of global and mesoscale meteorological model data to predict the transport and dispersion of tracer plumes over Washington, D.C. *Wea. Forecasting*, **21**, 383–394.
- Draxler, R., and J. L. Heffter, 1981: Workbook for estimating the climatology of regional-continental scale atmospheric dispersion over the United States. *NOAA Tech. Memo. ERL ARL-96*. NOAA, Air Resour. Lab. Silver Spring, Md.
- Draxler, R. R., and G. D. Hess, 1997: Description of the HYSPLIT_4 modeling system. *NOAA Tech. Memo. ERL ARL-224, Air Resources Laboratory, Silver Spring, MD*, 24 pp., NTIS PB98-116593.
- Draxler, R. R., and G. D. Hess, 1998: An overview of the HYSPLIT_4 modelling system for trajectories, dispersion, and deposition. *Aust. Meteor. Mag.*, **47**, 295–308.
- Draxler, R. R., and G. D. Rolph, 2012: Evaluation of the Transfer Coefficient Matrix (TCM) approach to model the atmospheric radionuclide air concentrations from Fukushima. *J. Geophys. Res.*, **117**, 10 pp, doi:10.1029/2011JD017205.
- Draxler, R., D. Arnold, S. Galmarini, M. Hort, A. Jones, S. Leadbetter, A. Malo, C. Maurer, G. Rolph, K. Saito, R. Servranckx, T. Shimbori, E. Solazzo, and G. Wotawa, 2013a: Evaluation of meteorological analyses for the radionuclide dispersion and deposition from the Fukushima Daiichi Nuclear Power Plant accident. *WMO Technical Publication*, **1120**, 64 pp. (available online at https://www.wmo.int/e-catalog/detail_en.php?PUB_ID=669)
- Draxler, R., P. Chen, M. Hort, A. Malo, K. Saito, and G. Wotawa, 2013b: World Meteorological Organization's evaluation of the radionuclide dispersion and deposition from the Fukushima Daiichi Nuclear Power Plant accident. *Proceeding, Special Symposium on the Transport and Diffusion of Contaminants from the Fukushima Dai-Ichi Nuclear Power Plant*. (available online at <https://ams.confex.com/ams/93Annual/webprogram/Paper215372.html>)
- Draxler, R., D. Arnold, M. Chino, S. Galmarini, M. Hort, A. Jones , S. Leadbetter, A. Malo, C. Maurer, G. Rolph, K. Saito, R. Servranckx, T. Shimbori, E. Solazzo, and G. Wotawa, 2015: World Meteorological Organization's model simulations of the radionuclide dispersion and deposition from the Fukushima Daiichi Nuclear Power Plant accident. *J. Environ. Radioact.*, **139**, 172–184.
- Fischer, B., 2012: Radionuclide emissions for the Fukushima Daiichi accident prepared for the UNSCEAR assessment. (private communication)

Fukushima Prefecture, 2011a:
http://www.pref.fukushima.lg.jp/sec_file/monitoring/m-3/20-50km0312-0331.pdf (in Japanese)

Fukushima Prefecture, 2011b:
http://www.pref.fukushima.lg.jp/sec_file/monitoring/m-1/7houbu0311-0331.pdf (in Japanese)

- Furuta, S., S. Sumiya, H. Watanabe, M. Nakano, K. Imaizumi, M. Takeyasu, A. Nakada, H. Fujita, T. Mizutani, M. Morisawa, Y. Kokubun, T. Kono, M. Nagaoka, H. Yokoyama, T. Hokama, T. Isozaki, M. Nemota, Y. Hiyama, T. Onuma, C. Kato, and T. Kurachi, 2011: Results of the Environmental Radiation Monitoring Following the Accident at the Fukushima Daiichi Nuclear Power Plant - Interim Report (Ambient Radiation Dose Rate, Radioactivity Concentration in the Air and Radioactivity Concentration in the Fallout), Radiation Protection Department, Nuclear Fuel Cycle Engineering Laboratories, Tokai Research and Development Center, Japan Atomic Energy Agency Tokai-mura, Naka-gun, Ibaraki-ken, *JAEA-Review* 2011-035, August, 89 pp. (in Japanese, available online at <http://jolissrch-inter.tokai-sc.jaea.go.jp/search/servlet/search?5031728>)
- Gauntt, R. O., R. K. Cole, C. M. Erickson, R. G. Gido, R. D. Gasser, S. B. Rodriguez, M. F. Young, S. Ashbaugh, M. Leonard, and A. Hill, 2001: MELCOR Computer Code Manuals, Vol. 3: Demonstration Problems, Version 1.8.5 May 2001, Sandia National Laboratories, SAND2001-0929P, Albuquerque, USA.
- Gifford, F. A., 1982: Horizontal diffusion in the atmosphere: A Lagrangian-dynamical theory. *Atmos. Environ.*, **16**, 505–512.
- Gifford, F. A., 1984: The random force theory: Application to meso and large-scale atmospheric diffusion. *Bound.-Layer Meteor.*, **30**, 159–174.
- Hayami, H., A. Sato, M. Tsuzaki, and H. Shimadera, 2012: Atmospheric transport and deposition modeling of radioactive materials released from the Fukushima Daiichi nuclear power plant. *CRIEPI report*, V11054. (in Japanese)
- Hertel, O., J. Christensen, E. H. Runge, W. A. H. Asman, R. Berkowicz, and M. F. Hovmand, 1995: Development and testing of a new variable scale air pollution model—ACDEP. *Atmos. Environ.*, **29**, 1267–1290.
- Hoffmann, W., R. Kebeasy, and P. Firbas, 2000: Introduction to the verification regime of the Comprehensive 5 Nuclear-Test-Ban Treaty. *Phys. Earth Planet. Inter.*, **113**, 5–9.
- Holtslag, A. A. M., and B. A. Boville, 1993: Local versus nonlocal boundary-layer diffusion in a global climate model. *J. Climate*, **6**, 1825–1842.
- Honda, Y., and K. Sawada, 2008: A new 4D-Var for mesoscale analysis at the Japan Meteorological Agency. *CAS/JSC WGNE Res. Act. Atmos. Ocea. Model.*, **38**, 01.07–01.08.
- Honda, Y., and K. Sawada, 2009: Upgrade of the operational mesoscale 4D-Var system at the Japan Meteorological Agency. *CAS/JSC WGNE Res. Act. Atmos. Ocea. Model.*, **39**, 01.11–01.12.
- Honda, Y., M. Nishijima, K. Koizumi, Y. Ohta, K. Tamiya, T. Kawabata, and T. Tsuyuki, 2005: A pre-operational variational data assimilation system for a non-hydrostatic model at the Japan Meteorological Agency: Formulation and preliminary results. *Q. J. Roy. Meteorol. Soc.*, **131**, 3465–3475.
- Hsu, S.-C., C.-A. Huh, C.-Y. Chan, S.-H. Lin, F.-J. Lin, and S. C. Liu, 2012: Hemispheric dispersion of radioactive plume laced with fission nuclides from the Fukushima nuclear event. *Geophys. Res.*

- Lett.*, **39**, L00G22, doi:10.1029/2011GL049986.
- Hu, X., D. Li, H. Huang, S. Shen, and E. Bou-Zeid, 2014: Modeling and sensitivity analysis of transport and deposition of radionuclides from the Fukushima Daiichi accident. *Atmos. Chem. Phys. Discuss.*, **14**, 2113–2173.
- Huijnen, V., J. Williams, M. van Weele, T. van Noije, M. Krol, F. Dentener, A. Segers, S. Houweling, W. Peters, J. de Laat, F. Boersma, P. Bergamaschi, P. van Velthoven, P. Le Sager, H. Eskes, F. Alkemade, R. Scheele, P. Nédélec, and H.-W. Pätz, 2010: The global chemistry transport model TM5: description and evaluation of the tropospheric chemistry version 3.0. *Geosci. Model Dev.*, **3**, 445–473.
- Hunt, B. R., E. J. Kostelich, and I. Szunyogh, 2007: Efficient data assimilation for spatiotemporal chaos: A local ensemble transform Kalman filter. *Physica D*, **230**, 112–126.
- Igarashi, Y., Y. Inomata, M. Aoyama, K. Hirose, H. Takahashi, Y. Shinoda, N. Sugimoto, A. Shimizu, and M. Chiba, 2009: Plausible change in Asian dust source suggested by atmospheric anthropogenic radionuclides-observation of single wet deposition events during spring of 2007. *Atmos. Environ.*, **43**, 2971–2980.
- Ikuta, Y., and Y. Honda, 2011: Development of 1D+4DVAR data assimilation of radar reflectivity in JNoVA. *CAS/JSC WGNE Res. Activ. Atmos. Ocea. Model.*, **41**, 01.09–01.10.
- Ishikawa, Y., 2010: Data assimilation of GPS precipitable water vapor into the JMA mesoscale numerical weather prediction model. *CAS/JSC WGNE Res. Act. Atmos. Ocea. Model.*, **40**, 01.13–01.14.
- Iwasaki, T., 2013: What happened from the accidental release of radioactive materials from the Fukushima Dai-Ichi Nuclear Power Plant on 3.11 in 2011? *Extended abstract, Special Symposium on the Transport and Diffusion of Contaminants from the Fukushima Dai-Ichi Nuclear Power Plant*, 4 pp. AMS annual meeting on 5th of January at Austin. (available online at <https://ams.confex.com/ams/93Annual/webprogram/Manuscript/Paper220977/AMS-Extended%20Abstract.Iwasaki.2013.pdf>)
- Iwasaki, T., T. Maki, and K. Katayama, 1998: Tracer transport model at Japan Meteorological Agency and its application to the ETEX data. *Atmos. Environ.*, **32**, 4285–4295.
- JCAC, 2011: <http://www.jcac.or.jp/site/senryo/taiki-kouka-back-list.html> (in Japanese)
- JMA, 2013: Outline of the operational numerical weather prediction at the Japan Meteorological Agency (March 2013), 188 pp. (available online at http://www.jma.go.jp/jma/jma-eng/jma-center/nwp/outline2013-nwp/pdf/outline2013_all.pdf)
- JMA, 2014: Specifications of the JMA's NWP models and Ensemble Prediction Systems. (available online at http://www.jma.go.jp/jma/jma-eng/jma-center/nwp/specifications_models.pdf)
- Jones, A. R., D. J. Thomson, M. C. Hort, and B. Devenish, 2007: The U.K. Met Office's next-generation atmospheric dispersion model, NAME III. In C. Borrego and A.-L. Norman, editors, Air Pollution and its Applications XVII. *Proceedings of the 27th NATO/CCMS International Technical Meeting on Air Pollution Modelling and its Application*, Springer,

580–589.

- Kajino, M., 2011: MADMS: Modal Aerosol Dynamics model for multiple Modes and fractal Shapes in the free-molecular and near-continuum regimes. *J. Aerosol Sci.*, **42**, 224–248.
- Kajino, M., and Y. Kondo, 2011: EMTACS: Development and regional-scale simulation of a size, chemical, mixing type, and soot shape resolved atmospheric particle model. *J. Geophys. Res.*, **116**, D02303, 28 pp, doi:10.1029/2010JD015030.
- Kajino, M., M. Deushi, T. Maki, N. Oshima, Y. Inomata, K. Sato, T. Ohizumi, and H. Ueda, 2012a: Modeling wet deposition and concentration of inorganics over Northeast Asia with MRI-PM/c. *Geosci. Model Dev.*, **5**, 1363–1375.
- Kajino, M., Y. Inomata, K. Sato, H. Ueda, Z. Han, J. An, G. Katata, M. Deushi, T. Maki, N. Oshima, J. Kurokawa, T. Ohara, A. Takami, and S. Hatakeyama, 2012b: Development of the RAQM2 aerosol chemical transport model and predictions of the Northeast Asian aerosol mass, size, chemistry, and mixing type. *Atmos. Chem. Phys.*, **12**, 11833–11856.
- Kaneyasu, N., H. Ohashi, F. Suzuki, T. Okada, and F. Ikemori, 2012: Sulfate aerosol as a potential transport medium of radio cesium from the Fukushima nuclear accident. *Environ. Sci. Technol.*, **46**, 5720–5726.
- Katata, G., M. Ota, H. Terada, M. Chino, and H. Nagai, 2012a. Atmospheric discharge and dispersion of radionuclides during the Fukushima Dai-ichi Nuclear Power Plant accident. Part I: source term estimation and local-scale atmospheric dispersion in early phase of the accident. *J. Environ. Radioact.*, **109**, 103–113.
- Katata, G., H. Terada, H. Nagai, and M. Chino, 2012b: Numerical reconstruction of high dose rate zones due to the Fukushima Daiichi Nuclear Power Plant accident. *J. Environ. Radioact.*, **111**, 2–12.
- Katata, G., M. Chino, T. Kobayashi, H. Terada, M. Ota, H. Nagai, M. Kajino, R. Draxler, M. C. Hort, A. Malo, T. Torii, and Y. Sanada, 2015: Detailed source term estimation of the atmospheric release for the Fukushima Daiichi Nuclear Power Station accident by coupling simulations of atmospheric dispersion model with improved deposition scheme and oceanic dispersion model. *Atmos. Chem. Phys.*, **15**, 1029–1070.
- Kawai, H., 2002: Forecast of sulfur dioxide flow from Miyake volcano with a high resolution regional transport model. *CAS/JSC WGNE Res. Act. Atmos. Ocea. Model.*, **32**, 05.24–05.25.
- Kitada, T., 1994: Modelling of transport, reaction and deposition of acid rain. *Kishou Kenkyu Note*, **182**, 95–117. (in Japanese)
- Kitada, T., G. R. Carmichael, and L. K. Peters 1986: Effects of dry deposition on the concentration-distribution of atmospheric pollutants within land- and sea-breeze circulations. *Atmos. Environ.*, **20**, 1999–2010.
- Kobayashi T., H. Nagai, M. Chino, and H. Kawamura, 2013: Source term estimation of atmospheric release due to the Fukushima Dai-ichi Nuclear Power Plant accident by atmospheric and oceanic dispersion simulations. *J. Nucl. Sci. Technol.*, **50**, 255–264.

- Kondo, H., T. Satomura, T. Takemura, H. Yamazawa, and A. Watanabe, 2012: A report of the special topic session “Current status and issues of numerical models for the transport of radioactive substances” at the 2011 Autumn Meeting. *Tenki*, **59**, 239–250. (in Japanese)
- Kondo, H., T. Yamada, M. Chino, T. Iwasaki, G. Katata, T. Maki, K. Saito, H. Terada, and H. Tsuruta, 2013: Report of the Special Symposium on the Transport and Diffusion of Contaminants from the Fukushima Dai-Ichi Nuclear Power Plant: Present Status and Future Directions. *Tenki*, **60**, 723–729. (in Japanese)
- Korsakissok, I., A. Mathieu, and D. Didier, 2013: Atmospheric dispersion and ground deposition induced by the Fukushima Nuclear Power Plant accident: a local-scale simulation and sensitivity study. *Atmos. Environ.*, **70**, 267–279.
- Kunii, M., 2013: Mesoscale data assimilation for a local severe rainfall event with the NHM-LETKF system. *Wea. Forecasting*, **29**, 1093–1105.
- Leadbetter, S., M. Hort, A. Jones, H. Webster, and R. Draxler, 2015: Sensitivity of the deposition of Caesium-137 from Fukushima Dai-ichi nuclear power plant on the wet deposition parameterisation in NAME. *J. Environ. Radioact.*, **139**, 200–211.
- Lim, K.-S. S., and S.-Y. Hong, 2010: Development of an effective double-moment cloud microphysics scheme with prognostic cloud condensation nuclei (CCN) for weather and climate models. *Mon. Wea. Rev.*, **138**, 1587–1612.
- Louis, J. F., M. Tiedtke, and J. F. Geleyn, 1982: A short history of the operational PBL parameterization at ECMWF. *Workshop on planetary boundary layer parameterization*. ECMWF, 59–79.
- Makihara, Y., 2000: Algorithms for precipitation nowcasting focused on detailed analysis using radar and raingauge data, Study on the Objective Forecasting Techniques. *Tech. Rep. MRI*, **39**, 63–111.
- Maryon, R. H., D. B. Ryall, and A. L. Malcolm, 1999: The NAME 4 dispersion model: Science documentation. *Technical report, UK Meteorological Office*. Turbulence Diffusion Note, **262**.
- Masson, O., A. Baeza, J. Bieringer, K. Brudecki, S. Bucci, M. Cappai, F. P. Carvalho, O. Connan, C. Cosma, A. Dalheimer, D. Didier, G. Depuydt, L. E. De Geer, A. De Vismes, L. Gini, F. Groppi, K. Guðnason, R. Gurriaran, D. Hainz, Ó. Halldórsson, D. Hammond, O. Hanley, K. Holeý, Zs. Homoki, A. Ioannidou, K. Isajenko, M. Jankovic, C. Katzlberger, M. Kettunen, R. Kierepko, R. Kontro, P. J. M. Kwakman, M. Lecomte, L. Leon Vintro, A.-P. Leppänen, B. Lind, G. Lujaniene, P. Mc Ginnity, C. Mc Mahon, H. Malá, S. Manenti, M. Manolopoulou, A. Mattila, A. Mauring, J. W. Mietelski, B. Møller, S. P. Nielsen, J. Nikolic, R. M. W. Overwater, S. E. Pálsson, C. Papastefanou, I. Penev, M. K. Pham, P. P. Povinec, H. Ramebäck, M. C. Reis, W. Ringer, A. Rodriguez, P. Rulík, P. R. J. Saey, V. Samsonov, C. Schlosser, G. Sgorbati, B. V. Silobritiene, C. Söderström, R. Sogni, L. Solier, M. Sonck, G. Steinhauser, T. Steinkopff, P. Steinmann, S. Stoulos, I. Sýkora, D. Todorovic, N. Tooloutalaie, L. Tositti, J. Tschiersch, A. Ugron, E. Vagena, A. Vargas, H. Wershofen, and O. Zhukova, 2011: Tracking of airborne radionuclides from the damaged Fukushima Dai-Ichi nuclear reactors by European networks. *Environ. Sci. Technol.*, **45**,

7670–7677.

- Mellor, G. L., and T. Yamada, 1974: A hierarchy of turbulence closure models for planetary boundary layers. *J. Atmos. Sci.*, **31**, 1791–1806.
- Mellor, G. L., and T. Yamada, 1982: Development of a turbulence closure model for geophysical fluid problems. *Rev. Geophys. Space Phys.*, **20**, 851–875.
- METI, 2011: <http://www.meti.go.jp/press/2011/06/20110603019/20110603019.html> (in Japanese)
- MEXT, 2007: SPEEDI System for Prediction of Environmental Emergency Dose Information, *Pamphlet of SPEEDI*.
- MEXT, 2011a: Monitoring information of environmental radioactivity level, Readings of dust sampling, <http://radioactivity.nsr.go.jp/en/list/200/list-1.html>
- MEXT, 2011b: Monitoring information of environmental radioactivity level, Readings of Environmental Radiation Level by emergency monitoring, <http://radioactivity.nsr.go.jp/en/list/207/list-1.html>
- MEXT, 2011c: Results of analyses of radionuclides (Cs-134 and Cs-137) in soil. (Retrieved from http://www.mext.go.jp/b_menu/shingi/chousa/gijyutu/017/shiryo/_icsFiles/afieldfile/2011/09/02/1310688_1.pdf)
- MEXT, 2011d: Results of airborne monitoring by the Ministry of Education, Culture, Sports, Science and Technology and the U.S. Department of Energy. Available online at
http://radioactivity.nsr.go.jp/ja/contents/4000/3710/24/1305820_20110506.pdf,
http://radioactivity.nsr.go.jp/ja/contents/5000/4858/24/1305819_0708.pdf,
http://radioactivity.nsr.go.jp/ja/contents/5000/4901/24/1910_1216.pdf
- Morino, Y., T. Ohara, and M. Nishizawa, 2011: Atmospheric behavior, deposition, and budget of radioactive materials from the Fukushima Daiichi nuclear power plant in March 2011. *Geophys. Res. Lett.*, **38**, doi:10.1029/2011GL048689.
- Morino, Y., T. Ohara, M. Watanabe, S. Hayashi, and M. Nishizawa, 2013: Episode analysis of deposition of radio cesium form the Fukushima Daiichi Nuclear Power Plant accident. *Environ. Sci. Technol.*, **47**, 2314–2322.
- Morrison, H., G. Thompson, and V. Tatarkii, 2009: Impact of cloud microphysics on the development of trailing stratiform precipitation in a simulated squall line: comparison of one- and two-moment schemes. *Mon. Wea. Rev.*, **137**, 991–1007.
- Nagai, H., M. Chino, and H. Yamazawa, 1999: Development of scheme for predicting atmospheric dispersion of radionuclides during nuclear emergency by using atmospheric dynamic model. *J. At. Energy. Soc. Japan*, **41**, 777–785. (in Japanese with English abstract)
- Nagata, K., 2011: Quantitative precipitation estimation and quantitative precipitation forecasting by the Japan Meteorological Agency. *RSMC Tokyo – Typhoon Center Technical Review*, **13**, 37–50. (available online at <http://www.jma.go.jp/jma/jma-eng/jma-center/rsmc-hp-pub-eg/techrev/text13-2.pdf>)
- NSC, 2010: The guideline for environmental radiation monitoring. (in Japanese, available online at

http://www.bousai.ne.jp/vis/shiryou/pdf/kankyou_monitor_h2204.pdf

- O'Brien, J. J., 1970: A note on the vertical structure of the eddy exchange coefficient in the planetary boundary layer. *J. Atmos. Sci.*, **27**, 1213–1215.
- Onogi, K., J. Tsutsui, H. Koide, M. Sakamoto, S. Kobayashi, H. Hatsushika, T. Matsumoto, N. Yamazaki, H. Kamahori, K. Takahashi, S. Kadokura, K. Wada, K. Kato, R. Oyama, T. Ose, N. Mannoji, and R. Taira, 2007: The JRA-25 Reanalysis. *J. Meteor. Soc. Japan*, **85**, 369–432.
- Park, S.-U., A. Choe, and M.-S. Park, 2013: Atmospheric dispersion and deposition of radionuclides (^{137}Cs and ^{131}I) released from the Fukushima Dai-ichi Nuclear Power Plant. *Comput. Water, Energy, Environ. Eng.*, **2**, 61–68.
- Pudykiewicz, J., 1989: Simulation of the Chernobyl dispersion with a 3-D hemispheric tracer model. *Tellus B*, **41**, 391–412.
- Ryall, D. B., and R. H. Maryon, 1998: Validation of the UK Met. Office's name model against the ETEX dataset. *Atmos. Environ.* **32**, 4265–4276.
- Saito, K., 2012: The Japan Meteorological Agency nonhydrostatic model and its applications to operation and research. *Atmospheric Model Applications*, 85–110, doi:10.5772/35368.
- Saito, K., T. Fujita, Y. Yamada, J. Ishida, Y. Kumagai, K. Aranami, S. Ohmori, R. Nagasawa, S. Kumagai, C. Muroi, T. Kato, H. Eito, and Y. Yamazaki, 2006: The operational JMA nonhydrostatic mesoscale model. *Mon. Wea. Rev.*, **134**, 1266–1298.
- Saito, K., J. Ishida, K. Aranami, T. Hara, T. Segawa, M. Narita, and Y. Honda, 2007: Nonhydrostatic atmospheric models and operational development at JMA. *J. Meteor. Soc. Japan*, **85B**, 271–304.
- Saito, K., T. Shimbori, and R. Draxler, 2013: JMA's regional ATM calculations for the WMO technical Task Team on meteorological analyses for Fukushima Daiichi Nuclear Power Plant accident. *Extended abstract, Special Symposium on the Transport and Diffusion of Contaminants from the Fukushima Dai-Ichi Nuclear Power Plant*, 5 pp. (available online at https://ams.confex.com/ams/93Annual/webprogram/Manuscript/Paper219086/AMS_ExtAbs_JM_ARATM.pdf)
- Saito, K., T. Shimbori, T. Hara, E. Toyoda, T. Kato, T. Fujita, K. Nagata, and Y. Honda, 2014: The WMO Technical Task Team on meteorological analyses for Fukushima Daiichi Nuclear Power Plant accident. *Sokko-jiho* (Japanese bulletin on operations at JMA), **81**, 1–30. (in Japanese, available online at <http://www.jma.go.jp/jma/kishou/books/sokkou/81/vol81p001.pdf>)
- Saito, K., T. Shimbori, and R. Draxler, 2015: JMA's regional atmospheric transport model calculations for the WMO technical task team on meteorological analyses for Fukushima Daiichi Nuclear Power Plant accident. *J. Environ. Radioact.*, **139**, 185–199.
- Sakamoto, M., 2013: Atmospheric Transport Model. Outline of the Operational Numerical Weather Prediction at the Japan Meteorological Agency, 95–98. (available online at http://www.jma.go.jp/jma/jma-eng/jma-center/nwp/outline2013-nwp/pdf/outline2013_03.pdf)
- Schöppner, M., W. Plastino, P. P. Povinec, G. Wotawa, F. Bella, A. Budano, M. De Vincenzi, and F. Ruggieri, 2011: Estimation of the time-dependent radioactive source-term from the Fukushima

- nuclear power plant accident using atmospheric transport modelling. *J. Environ. Radioact.*, **114**, 10–14.
- SCJ, 2014: A review of the model comparison of transportation and deposition of radioactive materials released to the environment as a result of the Fukushima Daiichi Nuclear Power Plant accident. *Report of working group for model intercomparison, Subcommittee to Review the Investigation on Environmental Contamination Caused by the Nuclear Accident, Committee on Comprehensive Synthetic Engineering, Science Council of Japan*. (available online at <http://www.scj.go.jp/ja/info/kohyo/pdf/kohyo-22-h140902-e1.pdf>)
- Seibert, P., D. Arnold, M. Gabriele, N. Arnold, K. Gufler, H. Kromp-Kolb, W. Kromp, P. Sutter, and A. Wenisch, 2012: Severe accidents of nuclear power plants in Europe: possible consequences and mapping of risk. *4th International Disaster and Risk Conference IDRC*, Davos, 26-30 August. (available online at <http://www.slideshare.net/GRFDavos/severe-accidents-of-nuclear-power-plants-in-europe-possible-consequences-and-mapping-of-risk>)
- Seino, N., H. Sasaki, J. Sato, and M. Chiba, 2004: High-resolution simulation of volcanic sulfur dioxide dispersion over the Miyake Island. *Atmos. Environ.*, **38**, 7073–7081.
- Sekiya, T. T., M. Kajino, and M. Kunii, 2013: Ensemble simulation of the atmospheric radionuclides discharged by the Fukushima nuclear accident. *Geophys. Res. Abstracts*, **15**, EGU2013-1695. (available online at <http://meetingorganizer.copernicus.org/EGU2013/EGU2013-1695.pdf>)
- Sekiya, T. T., M. Kunii, M. Kajino, and T. Shimbori, 2015: Horizontal resolution dependence of atmospheric simulations of the Fukushima nuclear accident using 15-km, 3-km, and 500-m grid models. *J. Meteor. Soc. Japan*, **93**, 49–64.
- Shao, Y., 2000: Physics and Modelling of Wind Erosion. *Kluwer Academic Pub.*, Dordrecht, pp. 268–270.
- Shimbori, T., Y. Aikawa, and N. Seino, 2009: Operational implementation of the tephra fall forecast with the JMA mesoscale tracer transport model. *CAS/JSC WGNE Res. Act. Atmos. Ocea. Model*, **39**, 05.29–05.30.
- Shimbori, T., Y. Aikawa, K. Fukui, A. Hashimoto, N. Seino, and H. Yamasato, 2010: Quantitative tephra fall prediction with the JMA mesoscale tracer transport model for volcanic ash: A case study of the eruption at Asama volcano in 2009. *Pap. Met. Geophys.*, **61**, 13–29, doi:10.2467/mripapers.61.13. (in Japanese with English abstract and figure captions)
- Shimbori T., R. Kai, Yo. Hayashi, Yu. Hayashi, A. Sugai, Y. Hasegawa, A. Hashimoto, A. Takagi, T. Yamamoto, and K. Fukui, 2014: Tephra fall predictions with the JMA regional atmospheric transport model: A case study of the eruptions at Shinmoe-dake volcano in 2011. *Pap. Met. Geophys.*, **65**, 75–107, doi:10.2467/mripapers.65.75. (in Japanese with English abstract and figure captions)
- Shoji, Y, 2009: A study of near real-time water vapor analysis using a nationwide dense GPS network

- of Japan. *J. Meteor. Soc. Japan*, **87**, 1–18.
- Skamarock, W. C., J. B. Klemp, J. Dudhia, D. O. Jill, D. M. Barker, M. G. Duda, X. Y. Huang, W. Wang, and J. G. Powers, 2008: A description of the advanced research WRF version 3. *Tech. Note, NCAR/TN-475+STR*, 125 pp., Natl. Cent. For Atmos. Res., Boulder, Colo.
- Smith, A. R., K. J. Thomas, E. B. Norman, D. L. Hurley, B. T. Lo, Y. D. Chan, P. V. Guillaumon, and B. G. Harvey, 2014: Measurements of fission products from the Fukushima Daiichi incident in San Francisco Bay Area air filters, automobile filters, rainwater, and food. *J. Environ. Protect.*, **5**, 207–221.
- Solazzo, E., and S. Galmarini, 2015: The Fukushima-Cs137deposition case study: properties of the Multi-Model ensemble. *J. Environ. Radioact.*, **139**, 226–233.
- Sportisse, B., 2007: A review of parameterizations for modelling dry deposition and scavenging of radionuclides. *Atmos. Environ.*, **41**, 2683–2698.
- Stohl, A., M. Hittenberger, and G. Wotawa, 1998: Validation of the Lagrangian particle dispersion model FLEXPART against large-scale tracer experiment data. *Atmos. Environ.* **32**, 4245–4264.
- Stohl, A., C. Forster, A. Frank, P. Seibert, and G. Wotawa, 2005, Technical note: The Lagrangian particle dispersion model FLEXPART version 6.2. *Atmos. Chem. Phys.*, **5**, 2461–2474.
- Stohl, A., P. Seibert, G. Wotawa, D. Arnold, J. F. Burkhart, S. Eckhardt, C. Tapia, A. Vargas, and T. J. Yasunari, 2012: Xenon-133 and caesium-137 releases into the atmosphere from the Fukushima Dai-ichi nuclear power plant: determination of the source term, atmospheric dispersion, and deposition. *Atmos. Chem. Phys.*, **12**, 2313–2343.
- Sugiyama, G., J. Nasstrom, B. Pobanz, K. Foster, M. Simpson, P. Vogt, F. Aluzzi, and S. Homann, 2012: Atmospheric dispersion modeling: challenges of the Fukushima Daiichi response. *Health Phys.*, **102**, 493–508.
- Suzuki, T., 1983: A theoretical model for dispersion of tephra: Arc Volcanism: Physics and Tectonics, D. Shimozuru and I. Yokoyama, Eds., *TERRAPUB*, Tokyo, pp. 95–113.
- Takano, I., Y. Aikawa, and S. Gotoh, 2007: Improvement of photochemical oxidant information by applying transport model to oxidant forecast. *CAS/JSC WGNE Res. Act. Atmos. Ocea. Model.*, **37**, 05.35–05.36.
- Takemura, T., H. Nakamura, M. Takigawa, H. Kondo, T. Satomura, T. Miyasaka, and T. Nakajima, 2011: A numerical simulation of global transport of atmospheric particles emitted from the Fukushima Daiichi Nuclear Power Plant. *SOLA*, **7**, 101–104.
- Takigawa, M., H. Nagai, Y. Morino, T. T. Sekiyama, H. Hayami, T. Tanaka, T. Nakajima, and T. Shibata, 2013: International intercomparison of atmospheric simulations of radionuclides. *Proceedings, the 54th Annual Meeting of Japan Society for Atmospheric Environment*, 136 p. (in Japanese)
- Tanaka, T. Y., and M. Chiba, 2005: Global simulation of dust aerosol with a chemical transport Model, MASINGAR. *J. Meteor. Soc. Japan*, **83A**, 255–278.
- Tanaka, T. Y., K. Orito, T. T. Sekiyama, K. Shibata, M. Chiba, and H. Tanaka, 2003: MASINGAR, a

- global tropospheric aerosol chemical transport model coupled with MRI/JMA98 GCM: Model description. *Pap. Met. Geophys.*, **53**, 119–138, doi:10.2467/mripapers.53.119.
- Tanaka, T. Y., M. Kajino, T. Maki, T. T. Sekiyama, Y. Igarashi, M. Chiba, and M. Mikami, 2012: Numerical simulation of atmospheric transport of radionuclides in Meteorological Research Institute. Japan Geoscience Union Meeting 2012.
- Terada, H., and M. Chino, 2008: Development of an atmospheric dispersion model for accidental discharge of radionuclides with the function of simultaneous prediction for multiple domains and its evaluation by application to the Chernobyl nuclear accident. *J. Nucl. Sci. Technol.*, **45**, 920–931.
- Terada, H., H. Nagai, A. Furuno, T. Kakefuda, T. Harayama, and M. Chino, 2008: Development of Worldwide Version of System for Prediction of Environmental Emergency Dose Information: WSPEEDI 2nd version. *Trans. At. Energy Soc. Japan*, **7**, 257–267. (in Japanese with English abstract)
- Terada, H., G. Katata, M. Chino, and H. Nagai, 2012: Atmospheric discharge and dispersion of radionuclides during the Fukushima Dai-ichi Nuclear Power Plant accident. Part II: verification of the source term and analysis of regional-scale atmospheric dispersion. *J. Environ. Radioact.*, **112**, 141–154.
- Torii, T., Y. Sanada, T. Sugita, A. Kondo, Y. Shikaze, M. Takahashi, M. Ishida, Y. Nishizawa, and Y. Urabe, 2012: Investigation of radionuclide distribution using aircraft for surrounding environmental survey from Fukushima Dai-ichi Nuclear Power Plant. *JAEA-Technologoy 2012-036*, Japan Atomic Energy Agency, 182 pp. (in Japanese with English abstract)
- Uliasz, M., 1990: Development of the mesoscale dispersion modeling system using personal computers. Part I: Models and computer implementation. *Z. Meteor.*, **40**, 110–120.
- UNSCEAR, 2013: Report of the United Nations Scientific Committee on the Effects of Atomic Radiation to the General Assembly. 17 pp. (available online at http://www.unscear.org/docs/reports/2013/13-85418_Report_2013_GA_Report.pdf)
- UNSCEAR, 2014: Levels and effects of radiation exposure due to the nuclear accident after the 2011 great east-Japan earthquake and tsunami. *UNSCEAR 2013 Report, Vol. 1, Scientific Annex A*. 321 pp. (available online at http://www.unscear.org/docs/reports/2013/13-85418_Report_2013_Annex_A.pdf)
- USDOE, 2011: US DOE/NNSA Response to 2011 Fukushima Incident, *United States Department of Energy*. (available online at <https://explore.data.gov/d/prrn-6s35>)
- Webster, H. N., and D. J. Thomson, 2011: Dry deposition modeling in a Lagrangian dispersion model. *Int. J. Environ. Pollut.*, **47**, 1–9.
- Webster, H. N., D. J. Thomson, B. T. Johnson, I. P. C. Heard, K. Turnbull, F. Marenco, N. I. Kristiansen, J. Dorsey, A. Minikin, B. Weinzierl, U. Schumann, R. S. J. Sparks, S. C. Loughlin, M. C. Hort, S. J. Leadbetter, B. J. Devenish, A. J. Manning, C. S. Witham, J. M. Haywood, and B. W. Golding, 2012: Operational prediction of ash concentrations in the distal volcanic cloud from

- the 2010 Eyjafjallajökull eruption. *J. Geophys. Res.*, **117**, D00U08, 17pp, doi:10.1029/2011JD016790.
- WGMSJ (Working Group on Emergency Response to Atmospheric Dispersion of Accidental Release of Radioactive Materials, Meteorological Society of Japan), 2014: Utilization of numerical prediction of atmospheric dispersion of accidentally released radioactive materials (in Japanese, available online at <http://www.metsoc.jp/default/wp-content/uploads/2014/12/teigen-201412.pdf>)
- Wilson, J. D., F. J. Ferrandino, and G. W. Thurtell, 1989: A relationship between deposition velocity and trajectory reflection probability for use in stochastic Lagrangian dispersion models. *Agric. For. Meteor.*, **47**, 139–154.
- Winiarek, V., M. Bocquet, O. Saunier, and A. Mathieu, 2012: Estimation of errors in the inverse modeling of accidental release of atmospheric pollutant: Application to the reconstruction of the cesium-137 and iodine-131 source terms from the Fukushima Daiichi power plant. *J. Geophys. Res.*, **117**, D05122, 16 pp, doi:10.1029/2011JD016932.
- Winiarek, V., M. Bocquet, N. Duhanyan, Y. Roustan, O. Saunier, and A. Mathieu, 2014: Estimation of the caesium-137 source term from the Fukushima Daiichi nuclear power plant using a consistent joint assimilation of air concentration and deposition observations. *Atmos. Environ.*, **82**, 268–279.
- WMO, 2010: Manual on the Global Data-processing and Forecasting System. WMO, **485**.
- WMO, 2011, Final report of the Meeting of the WMO Task Team on Meteorological Analyses for Fukushima Daiichi Nuclear Power Plant Accident, Geneva, Switzerland, 30 November – 2 December 2011. 27 pp. (available online at http://www.wmo.int/pages/prog/www/CBS-Reports/documents/FinalRep_TT_FDnpp_v6.pdf)
- WMO, 2012a, Final report of the Second Meeting of the WMO Technical Task Team on Meteorological Analyses for Fukushima-Daiichi Nuclear Power Plant Accident, London, UK , 1 – 3 May 2012. 13 pp. (available online at http://www.wmo.int/pages/prog/www/CBS-Reports/documents/FinalReport_TTMetAnalyFDnpp.pdf)
- WMO, 2012b, Final report of the Third Meeting of the WMO Technical Task Team on Meteorological Analyses for Fukushima-Daiichi Nuclear Power Plant Accident, Vienna, Austria , 3 – 5 December 2012. 76 pp. (available online at <http://www.wmo.int/pages/prog/www/CBS-Reports/documents/FINAL-REPORT-Vienna-Dec2012.pdf> and http://www.wmo.int/pages/prog/www/CBS-Reports/documents/WMO_fnpp_final_AnnexIII_4Fe_b2013_REVISED_17June2013.pdf)
- Wotawa, G., L.-E. De Geer, P. Denier, M. Kalinowski, H. Toivonen, R. D'Amours, F. Desiato, J.-P. Issartel, M. Langer, P. Seibert, A. Frank, C. Sloan, and H. Yamazawa, 2002: Atmospheric transport modeling in support of CTBT verification—overview and basic concepts. *Atmos. Environ.*, doi:10.1016/S1352-2310(03)00154-7.
- Wotawa, G., R. Draxler, D. Arnold, S. Galmarini, M. Hort, A. Jones, S. Leadbetter, A. Malo, C.

- Maurer, G. Rolph, K. Saito, R. Servranckx, T. Shimbori, and E. Solazzo, 2013: Transport and deposition of radionuclides after the Fukushima nuclear accident: international model inter-comparison in the framework of a WMO Task Team. *Geophys. Res. Abstracts*, **15**, EGU2013-3193. (available online at <http://meetingorganizer.copernicus.org/EGU2013/EGU2013-3193.pdf>)
- Yamazawa, H., A. Furuno, and M. Chino, 1998: Evaluation of a long-range Lagrangian dispersion model with ETEX. *Atmos. Environ.*, **32**, 4343–4349.
- Yasunari, T. J., A. Stohl, R. S. Hayano, J. F. Burkhardt, S. Eckhardt, and T. Yasunari, 2011: Cesium-137 deposition and contamination of Japanese soils due to the Fukushima nuclear accident. *Proc. Natl. Acad. Sci.*, **108**, 19530–19534.
- Yukimoto, S., H. Yoshimura, M. Hosaka, T. Sakami, H. Tsujino, M. Hirabara, T. Y. Tanaka, M. Deushi, A. Obata, H. Nakano, Y. Adachi, E. Shindo, S. Yabu, T. Ose, and A. Kitoh, 2011: Meteorological Research Institute-Earth System Model Version 1 (MRI-ESM1) —Model Description—. *Tech. Rep. MRI*, **64**, 88 pp, doi:10.11483/mritechrepo.64.
- Yukimoto, S., Y. Adachi, M. Hosaka, T. Sakami, H. Yoshimura, M. Hirabara, T. Y. Tanaka, E. Shindo, H. Tsujino, M. Deushi, R. Mizuta, S. Yabu, A. Obata, H. Nakano, T. Koshiro, T. Ose, and A. Kitoh, 2012: A New Global Climate Model of the Meteorological Research Institute: MRI-CGCM3 —Model Description and Basic Performance—. *J. Meterol. Soc. Japan*, **90A**, 23–64.